

Antimatter in the Milky Way

C. Bambi^{a,b,*} and A.D. Dolgov^{a,b,c,†}

February 5, 2008

^aIstituto Nazionale di Fisica Nucleare, Sezione di Ferrara, I-44100 Ferrara, Italy

^bDipartimento di Fisica, Università degli Studi di Ferrara, I-44100 Ferrara, Italy

^cInstitute of Theoretical and Experimental Physics, 113259, Moscow, Russia

Abstract

Observational signatures of existence of antimatter objects in the Galaxy are discussed. We focus on point-like sources of gamma radiation, diffuse galactic gamma ray background and anti-nuclei in cosmic rays.

*E-mail: bambi@fe.infn.it

†E-mail: dolgov@fe.infn.it

1 Introduction

One can conclude from simple considerations that there is much more matter than antimatter around us [1, 2]. The Earth and the Solar System are evidently made of matter and the very small antiproton-to-proton ratio in cosmic rays, $\sim 10^{-4}$, suggests a secondary origin of antiprotons by cosmic ray collisions in the interstellar medium and an absence of a large amount of antimatter in the Galaxy. Still an excess of cosmic antiprotons at low energies [3] might point to non-standard sources of their production and, in particular, to some antimatter objects in our neighborhood. A similar conclusion may be obtained from observations of gamma rays originating from e^+e^- annihilation [4, 5]. Though a conventional mechanism of positron production is the most probable one, light dark matter might be a possible source of positrons, especially because of the observation of the 511 keV line from the halo emission, see the above quoted papers. This explanation suffers from a necessity to fine-tune the mass of the dark matter particles, so that they would decay or annihilate into non-relativistic positrons. Another possible source of positrons could be primordial antimatter in the Galaxy. For other galaxies the sensitivity is not high enough to exclude or observe significant $\bar{p}p$ and e^+e^- annihilation indicating possible cosmic antimatter objects, but galaxies dominated by antimatter are excluded in our galactic cluster, i.e. up to distances of about 10 Mpc [1], by non-detection of γ rays which would be produced by the annihilation of the galactic antimatter with the matter from the infalling intergalactic gas.

On the other hand, *a priori* we could expect an approximately charge symmetric universe, or at least a universe with some considerable amount of antimatter, since matter and antimatter have (almost) the same properties.

Cosmological matter-antimatter asymmetry can be expressed in terms of the baryon-to-photon ratio, β . The Big Bang Nucleosynthesis (BBN) [6] and the Cosmic Microwave Background Radiation (CMBR) [7] provide two independent measurements which are consistent with each other

$$\beta = \frac{n_B - n_{\bar{B}}}{n_\gamma} \approx 6 \cdot 10^{-10} \quad (1)$$

where $n_B \gg n_{\bar{B}}$, whereas the freeze-out abundances in a homogeneous baryo-symmetric universe would be $n_B/n_\gamma = n_{\bar{B}}/n_\gamma \sim 10^{-18}$ [8].

There can be three possible types of cosmological matter-antimatter asymmetry:

1. β is constant and the universe is 100% matter dominated.
2. The universe is globally baryo-symmetric. It consists of equal amount of similar domains of matter and antimatter.
3. The universe has a non-vanishing average baryonic charge, but β is not spatially constant and can even be negative in some space regions. In other words there could be lumps of antimatter in matter dominated universe.

The type of the asymmetry depends upon the mechanism of baryogenesis which took place in the early universe. The baryogenesis scenarios mostly focus on the first possibility, for review see Refs. [9]-[12], but at present there is neither experimental nor observational evidence in favor of one model over another, since the involved physics operated at such high energies that it is difficult or impossible to test it in laboratories on the Earth. On the other hand, a globally baryo-symmetric universe is certainly theoretically appealing, but it

seems observationally excluded or, to be more precise, the size of the domain where we live should be (at least) comparable to the present day cosmological horizon [13].

Most interesting phenomenologically is the third case because it allows for existence of anti-objects in our neighborhood and hence for peculiar features which may be observed in a near future, thanks to the advent of antimatter research projects such as AMS and PAMELA. A small amount of antimatter is not excluded even nearby in the Galaxy. The aim of our paper is to consider phenomenological manifestation of that. The effects from antimatter objects in the Galaxy were analyzed also in Refs. [14] for a different mechanism of antimatter creation and because of that for restricted types of such objects.

The content of the paper is the following. In Sec. 2 we briefly describe a mechanism for generation of lumps of antimatter in baryon dominated universe. Their cosmological evolution is considered in Sec. 3 and matter-antimatter annihilation in contemporary universe is discussed in Sec. 4. In Sec. 5 we focus on point-like sources of γ rays, in Sec. 6 on the diffuse galactic gamma ray background, and in Sec. 7 on the possibility of anti-nuclei in cosmic rays. In Sec. 8 we speculate on more violent events, where large amounts of matter and antimatter come into contact. The results are summarized in Sec. 9.

2 Antimatter in baryon asymmetric universe

Baryon asymmetric universe with high density regions of matter and antimatter could be created if there existed two different sources of CP violation [15]: the background baryon asymmetry could be provided by an explicit violation of CP in the Lagrangian, whereas small bubbles with very high baryon asymmetry could be produced by the presence of a stochastic or dynamical violation of CP. The concrete models are considered e.g. in Refs. [15, 16].

In what follows we will use as a reference the scenario proposed in paper [17]. The starting point is the Affleck-Dine mechanism [18], where scalar fields with non-zero baryonic and/or leptonic charges (predicted in supersymmetric extensions of the Standard Model) have the potential with flat directions, that is the directions along which the potential energy does not change. As a toy-model, we can consider the potential of the form:

$$U(\chi) = \lambda|\chi|^4(1 - \cos 4\theta) , \quad (2)$$

where $\chi = |\chi|e^{i\theta}$ is the scalar field with baryonic charge $B \neq 0$. Potential (2) has four flat directions along $\cos 4\theta = 1$. It is not invariant under the transformation $\chi \rightarrow \chi e^{i\alpha}$, i.e. B is not conserved. Due to the infrared instability of massless (or light, $m_\chi \ll H$) fields in de Sitter space-time [19] during inflation, χ can condense along the flat directions of the potential, acquiring a large expectation value. In the course of the cosmological expansion the Hubble parameter drops down and when the mass of the field, m_χ , exceeds the universe expansion rate H , χ evolves to the equilibrium point $\chi = 0$ and the baryonic charge stored in the condensate is transformed into quarks by B -conserving processes. Since here CP is violated stochastically by a chaotic phase of the field χ , then during the motion to the equilibrium state the matter and antimatter domains in a globally symmetric universe would be created. An interesting feature of the model is that regions with a very high β , even close to one, could be formed.

If the scalar field χ is coupled to the inflaton Φ with an interaction term of the kind

$$V(\chi, \Phi) = \lambda|\chi|^2(\Phi - \Phi_1)^2 , \quad (3)$$

the “gates” to the flat directions might be open only for a short time when the inflaton field Φ was close to Φ_1 . In this case, the probability of the penetration to the flat directions is small and χ could acquire a large expectation value only in a tiny fraction of space. The universe would have a homogeneous background of baryon asymmetry $\beta \sim 6 \cdot 10^{-10}$ generated by the same field χ which did not penetrate to larger distance through the narrow gate or by another mechanism of baryogenesis, while the high-density matter, $\beta > 0$, and antimatter, $\beta < 0$, regions would be rare, while their contribution to the cosmological mass density might be significant or even dominant. Let us call these bubbles with high baryonic number density B-balls.

Since the conditions for B-ball creation had been prepared during inflation, their initial radius, R_B , might quite naturally exceed the cosmological horizon.

In a simple model, the mass spectrum of B-balls has a log-normal form [17]

$$\frac{dN}{dM} = C \exp \left[-\gamma \ln^2 \left(\frac{M}{M_0} \right) \right], \quad (4)$$

where C , γ and M_0 are unknown parameters of the underlying theory. Depending on their values and on β , which are stochastic quantities, such bubbles could form clouds of matter or antimatter with high (anti)baryon number density, more compact object like (anti)stars, and primordial black holes. If all antimatter is hidden inside anti-black holes it would be unobservable. However, it is natural to expect that there could be abundant anti-stars (either normal or compact ones, as e.g. white dwarfs) or clouds of antimatter with higher than normal baryon density. Such antimatter objects could survive in the early universe due to their higher density, invalidating the bounds of Ref. [14].

The compact matter/antimatter objects created by the described mechanism might make a part of, or even the whole, cosmological dark matter¹. An interesting feature of such dark matter is that it consists of (stellar size) “particles” with dispersed masses. In particular, even if M_0 is close to the solar mass, there is a non-zero probability that on the tail of distribution (4) very heavy black holes of millions solar masses might be created. In particular, if there is one heavy black hole per galaxy, they could serve as seeds for galaxy formation. On the other hand, in the non-collapsed regions with high baryonic number density primordial nucleosynthesis proceeded with large β , producing nuclei heavier than those formed in the standard BBN [21]. If these regions are in our neighborhood, an observation of heavy anti-nuclei in cosmic rays would be plausible.

A different model of creation of (much smaller size) compact anti matter objects was suggested in Ref. [22]. Their observational signatures in cosmic gamma ray radiation are analyzed in Ref. [23].

3 Cosmological evolution of B-balls

When baryogenesis was over, the regions with high values of $|\beta|$ had the same energy density as the background. Only the chemical content was different. These are the so called isocurvature density perturbations. On the boundary of the bubbles the density contrast should be non-zero due to the gradient term, $|\partial\chi|^2$, but it is relatively insignificant. Moreover, this term disappeared at later stage when χ relaxed down to the equilibrium

¹According to Ref. [20], the present observational data allow (at the 95% C.L.) macroscopic compact objects with masses greater than $10^{-2} M_\odot$ to constitute 88% or less of the cosmological dark matter, so 100% is not highly probable but still allowed.

point, $\chi = 0$. After the electroweak phase transition, when quarks acquired masses due to the Higgs field condensate, the density contrast became nonzero, especially because of the large mass of t -quark, $m_t \sim 150$ GeV. A short life-time of t and other heavy quarks and a large number of different quark species in the primeval plasma make the density contrast relatively small and we ignore it in what follows.

More essential is the QCD phase transition when quarks became confined forming non-relativistic nucleons. The density contrast of B-bubbles would be equal to:

$$r_B = \frac{\delta\rho}{\rho} = \frac{\beta n_\gamma m_N}{\pi^2 g_* T^4/30} = \beta \frac{0.7 m_N}{g_* T} \approx 0.068 \beta \left(\frac{10.75}{g_*} \right) \left(\frac{m_N}{T} \right), \quad (5)$$

where $m_N \approx 940$ MeV is the nucleon mass, T is the temperature of the cosmic plasma, and $g_* \approx 10$ is the number of particle species “living” in the plasma. The nonrelativistic baryonic matter starts to dominate inside the bubble at

$$T = T_{in} \approx 65 \beta \text{ MeV}. \quad (6)$$

The mass inside a baryon-rich bubble at the radiation dominated stage is

$$M_B = \frac{4\pi}{3} R_B^3 \rho = (1 + r_B) M_{Pl}^2 t \left(\frac{R_B}{2t} \right)^3 \approx 2 \cdot 10^5 M_\odot (1 + r_B) \left(\frac{R_B}{2t} \right)^3 \left(\frac{t}{\text{sec}} \right), \quad (7)$$

where $M_\odot$ is the solar mass and we assumed $\rho = 3M_{Pl}^2(1 + r_B)/32\pi t^2$. If at the moment of the horizon crossing, i.e. at $R_B = 2t$, the density contrast would be large, $r_B \geq 1$, the bubble gravitational radius R_g would be larger than R_B and the bubble should form a primordial black hole (BH). The BH masses are determined by the plasma temperature at the moment of horizon crossing. Assuming the approximate relation $t/\text{sec} \approx (\text{MeV}/T)^2$, we find $M_{BH} \approx 4 \cdot 10^5 M_\odot (t/\text{sec})$. The cosmological QCD phase transition took place at $t = 10^{-5} - 10^{-4}$ sec and the masses of the first formed BHs should be near 10 solar masses. The baryon-rich bubbles with larger radius could make much heavier BHs.

Smaller bubbles with $R_B < 2t$ and $r_B \sim 1$ could form compact stellar type objects with masses given by Eq. (7) and with the initial (that is at $T = T_{in}$) mass density of non-relativistic matter:

$$\rho_B \approx 4.4 \cdot 10^5 r_B \left(\frac{\text{sec}}{t_{in}} \right)^2 \text{ g/cm}^3 \approx 10^{13} \beta^4 \text{ g/cm}^3, \quad (8)$$

where Eq. (6) and $t/\text{sec} \approx (\text{MeV}/T)^2$ were used.

The subsequent evolution of B-balls depends upon the ratio of their mass to the Jeans mass. The Jeans wave length is:

$$\lambda_J = c_s \sqrt{\frac{\pi M_{Pl}^2}{\rho}} = t \sqrt{\frac{32\pi^2}{3} \frac{T}{m_N}} \approx 10t \sqrt{T/m_N} \quad (9)$$

where the speed of sound is $c_s \approx \sqrt{T/m_N}$ and $\rho = 3m_{Pl}^2/32\pi t^2$. It is evident from this expression that λ_J exceeds horizon, $l_h = 2t$, if $T < 40$ MeV. If we take the time and temperature in Eq. (9) at the beginning of matter dominance in the B-bubble given by Eq. (6), the initial value of λ_J would be

$$\lambda_J^{(in)} \approx 6.2 \cdot 10^{-4} \beta^{-3/2} \text{ s}. \quad (10)$$

Correspondingly the initial value of the Jeans mass is

$$M_J \approx 135 (T/m_N)^{3/2} M_{Pl}^2 t \approx 100 M_\odot \beta^{-1/2}. \quad (11)$$

For relativistic expansion the temperature drops as $T \sim 1/a$, where $a(t)$ is the cosmological scale factor. In such a case the Jeans mass would rise with time. However, the temperature of matter dominated B-bubbles drops much faster, $T \sim 1/a^2$, and the Jeans mass would decrease with time in agreement with intuitive expectations. According to Eq. (11) Jeans mass scales as $M_J \sim T^{3/2} t$. For nonrelativistic expansion law, $a \sim t^{2/3}$ and consequently

$$M_J \sim 100 M_\odot (a_{in}/a)^{3/2}, \quad (12)$$

where a_{in} is cosmological scale factor at the onset of matter dominance inside B-ball.

M_J became of the order of the solar mass for $a/a_{in} \sim 20$. For an estimate let us take $T_{in} \sim 10$ MeV. According to Eq. (6), it is realized for $\beta \approx 0.15$. Correspondingly the temperature inside B-bubble when M_J dropped down to $M_\odot$, would be

$$T_{in}(a_{in}/a)^2 \approx 0.025 \text{ MeV}. \quad (13)$$

The mass density evolved as $1/a^3$ and thus it became

$$\rho_B = \rho_B^{(in)} (a_{in}/a)^3 \approx 6 \cdot 10^5 \text{ g/cm}^3 \quad (14)$$

to the moment when $M_J = M_\odot$. Correspondingly the radius of such B-ball (i.e. the Jeans wave length) would be $R_B \sim 10^9$ cm.

It is noteworthy that B-bubbles would be supported against expansion by the pressure of the hotter surrounding relativistic plasma. Since the photon mean free path inside B-bubbles is quite small, see below, the heat exchange takes place only on the surface and is not efficient. The external pressure slows down the expansion of B-balls and their adiabatic cooling.

If $\lambda_J < R_B$, the baryon-rich bubbles with such radius would decouple from the cosmological expansion and for $R_B < 2t$ they would not be black holes but compact star-like objects. According to Eq. (14) their mass density could be much larger than the density of the normal stars and similar to that of compact stars. They might survive to the present time or, if nuclear reactions inside would be significant, they would explode enriching outer space with heavy elements or anti-elements. Since their temperature at the moment of formation is very high, the pressure is also high, higher than the usual pressure in normal stars powered by nuclear reactions. So we assume that such objects mostly survive to the present time. After a while, their temperature would become larger than the cosmological temperature because they stopped expanding and the temperature is not red-shifted. They would cool down because of radiation from the surface into external colder space. Their luminosity can be estimated as $L \sim T_{surf}^4 R_B^2$, where T_{surf} is the surface temperature. With infinitely large thermal conductivity, the life-time with respect to the cooling would be quite short $\tau \sim T^4 R_B^3 / L \sim R_B$. Of course this result is an underestimate and the real cooling time must be much longer. Without additional sources of energy, the cooling time should be of the order of the photon diffusion time, t_{diff} , inside these compact objects. If the photon mean free path l_{free} is smaller than the radius R_B the diffusion time from the center to the surface can be estimated as

$$t_{diff} \sim R_B^2 / \tau_{free}. \quad (15)$$

The mean free time of photons is equal to

$$\tau_{free} \sim 1/(\sigma_{e\gamma} n_e), \quad (16)$$

where n_e is the number density of electrons and $\sigma_{e\gamma}$ is the photon-electron cross-section. For low energy photons, $E_\gamma < m_e$, it is equal to the Thomson cross-section, $\sigma_{Th} = 8\pi\alpha^2/3m_e^2$. For large energies it drops as inverse center-of-mass energy squared.

The photon diffusion is quite slow and the cooling time of baryon-rich compact objects is determined by t_{diff} . Assuming that $\sigma_{e\gamma} = \sigma_{Th}$ we obtain:

$$t_{diff} \approx 2 \cdot 10^{11} \text{ sec} \left(\frac{M_B}{M_\odot} \right) \left(\frac{\text{sec}}{R_B} \right) \left(\frac{\sigma_{e\gamma}}{\sigma_{Th}} \right). \quad (17)$$

The mass and size of the baryonic bubbles depends upon the parameters of the model of their creation. We assume, for the sake of estimate, that they predominantly have mass close to the solar one and use the results presented after Eq. (12). The thermal energy of a solar mass B-bubble taken at the moment when the Jeans mass dropped down to $M_\odot$ is determined by the thermal energy of nucleons and electrons, $E_{th} = 3T$. Taking $T = 25$ keV, see Eq. (13), we find for the total energy stored inside the B-ball

$$E_{therm}^{(tot)} = 3TM_B/m_N \approx 1.5 \cdot 10^{29} \text{ g} \approx 1.5 \cdot 10^{50} \text{ erg}. \quad (18)$$

With the cooling time given by Eq. (17) the luminosity of B-bubble would be $L \approx 10^{39}$ erg/sec, i.e. 10^6 of the solar luminosity. Such B-bubbles would be quite bright sources of keV radiation. However, they would stop radiating their thermal before hydrogen recombination. The temperature of CMBR at $t = t_{diff} \sim 10^{11}$ s, would be about 3 eV. If the B-bubbles make all dark energy, i.e. $\Omega_{BB} = 0.25$, their mass density at t_{diff} would be equal to that of CMBR. Correspondingly the energy density of the emitted keV radiation would be at most $10^{-4} - 10^{-5}$ of CMBR. This radiation would red-shift today by 10^{-4} and move to eV region, i.e. to background light.

More efficient source of power could be nuclear reactions that might ignite inside B-balls. In the considered example with $\rho \sim 10^6$ g/cm³ B-ball has the properties similar to those of the core of red giant at the initial stage of its evolution. The main source of energy under these conditions would be helium-4 burning, $3\text{ }^4\text{He} \rightarrow \text{}^{12}\text{C}$. However, the temperature would be somewhat larger, $T \sim 25$ KeV, instead of 10 keV. Since the probability of the above reaction is proportional to T^{40} , the life-time of such helium flash would be extremely short. Naively taken these numbers, we obtain life-time about 10 s. However, this simple estimate can be wrong by several orders of magnitude because the efficiency of the process is very much different from that in normal giant star. Still helium would be burnt very quickly and later nuclear reactions would be presumably insignificant. More accurate estimates would demand development of astrophysics for such unusual objects as B-balls and this is not a subject of this work. In the following sections we consider observational manifestations of different types of B-balls allowing for existence of any type of such objects. Here we note only that if the nuclear physics of B-balls with the chosen above parameters is similar to that of the giant star core, the result of helium flash would be a compact star with mass density of about 10^4 g/cm³. The flash should take place at relatively early stage of cosmological expansion to be safe from the CMBR restrictions. Indeed that relative energy density of B-balls would be $\rho_{BB}/\rho_\gamma < 10^4/z$. Since nuclear reactions result in emission of approximately 10^{-3} of the total mas of a star, and we do not allow than more 10^{-4} distortion of CMBR, we

conclude that such B-balls should consume their helium at cosmological red-shift $z > 10^5$, i.e. at $T \approx 20$ eV and $t < 3 \cdot 10^9$ s, which does not look unreasonable in view of the presented above estimate of the life-time ~ 10 s.

KeV photons coming from the cooling of B-balls would heat up cosmic electrons and they in turn might distort the spectrum of CMBR. However, since the total energy influx is about $10^{-4} - 10^{-5}$ of the total energy density of CMBR, the spectrum distortion should be not more than at the same level. More detailed calculations are in order.

If B-bubbles consist of antimatter, the annihilation of the background matter on their surface would create an additional radiation. However, due to a strong coupling of protons and electrons to photons before hydrogen recombination, the proton diffusion to an anti-B-bubble is quite slow and it is easy to see that the process is not efficient. After recombination the number of annihilations on the surface of one B-ball per unit time would be

$$\dot{N} = 4\pi R_B^2 \beta n_\gamma V_p \approx 10^{31} V_p \left(\frac{T}{0.1 \text{ eV}} \right)^3 \left(\frac{R_B}{10^9 \text{ cm}} \right)^2, \quad (19)$$

where $\beta = n_B/n_\gamma = 6 \cdot 10^{-10}$ and $V_p \sim 10^{-5}$ is the hydrogen velocity. Assuming that the surface annihilation lasted approximately for one the Hubble time at $T \sim 0.1$ eV, i.e. $t = t_H \approx 10^{13}$ s and that the mass density of B-bubbles is not larger than $10 \rho_{CMBR}$, we obtain that the energy density of gamma radiation from the annihilation on the surface is smaller than 10^{-16} of that of CMBR.

A more detailed analysis is of course necessary but it does not fit the frames of this letter. The dynamics and astrophysics of B-bubbles in the early universe will be considered elsewhere. There could be quite interesting signatures of B-bubbles as e.g. gamma ray background and background light. They might also have an observable impact on CMBR. We will not go into these interesting phenomena here. We only want to demonstrate that the model of Ref. [17] allows an early production of dark stellar-mass objects consisting of matter and antimatter which may survive to the present time. Such objects would behave as the usual cold dark matter and so they would populate galactic halos in contrast to the normal stars which live in the smaller luminous parts of galaxies.

4 Antimatter in contemporary universe

In what follows we will not dwell on possible scenarios of antimatter creation but simply consider phenomenological consequences of their existence in the present day universe, in particular in the Galaxy, allowing for any types of such objects. According to the discussion in the previous section, astronomically interesting domains of antimatter probably populate galactic halos but we will not confine ourselves to this case only, but consider phenomenology of more general situation. If high density regions of antimatter have survived up to the present time, we assume that astronomical anti-objects can be either inside and/or in the halo of our galaxy. In particular there could be anti-clouds, anti-stars, anti-stellar clusters and anti-black holes (i.e. black holes generated by the gravitational collapse of antimatter; they may be distinguishable from “standard” black holes if they were surrounded by an anti-atmosphere). Of course, the presence of anti-objects in the Galaxy today should lead to the production of the gamma radiation from matter-antimatter annihilation. Hence we would expect ~ 100 MeV γ from the decay of π^0 -mesons produced in $p\bar{p}$ -annihilation, with an average of four γ per annihilation, and 2γ from e^+e^- -annihilation with $E = 0.511$ MeV if e^+e^- annihilate at rest. In addition to the slow background positrons there should be also

energetic secondary positrons produced by pion decays from $\bar{p}p$ -annihilation. Astronomical observations are seemingly more sensitive to $\bar{p}p$ -annihilation because the total energy release in $\bar{p}p$ -annihilation is 3 orders of magnitude larger than that in e^+e^- -annihilation and the galactic gamma ray background at 100 MeV is several orders of magnitude lower than the one at 0.5 MeV. On the other hand, e^+e^- -annihilation gives the well defined line which is easy to identify.

In order to study the observational constraints on the number of such anti-objects and to speculate on possible future investigations, it is helpful to distinguish two different regimes of matter-antimatter annihilation, depending on the ratio R/λ_{free} , where R is the size of the anti-object and

$$\lambda_{free} = 1/(\sigma_{ann} n_{\bar{p}}) \quad (20)$$

is the proton or electron mean free path inside an anti-object, with σ_{ann} and $n_{\bar{p}}$ being respectively the annihilation cross section for $\bar{p}p$ or e^+e^- (they have similar order of magnitude) and the antiproton number density in the anti-object.

4.1 Volume annihilation

Volume annihilation takes place if λ_{free} is larger than the radius R_B , that is:

$$R_B \sigma_{ann} n_{\bar{p}} \approx 3 \cdot 10^{-2} \left(\frac{R_B}{0.1 \text{ pc}} \right) \left(\frac{n_{\bar{p}}}{10^4 \text{ cm}^{-3}} \right) \left(\frac{\sigma}{10^{-23} \text{ cm}^2} \right) < 1. \quad (21)$$

One should remember that $\sigma_{ann} \sim 1/v$ where v is the velocity of the annihilating particles in their center-of-mass frame and thus the mean-free path is different for protons with different velocities. We assume that the typical velocities are of the order of $10^{-3}c$. The effective time of annihilation, $\tau = \lambda_{free}/v$ does not depend on velocity of the annihilating particles.

The annihilation rate per unit time and volume is equal to

$$\dot{n}_{\bar{p}} = \sigma_{ann} v n_p n_{\bar{p}}, \quad (22)$$

where $n_p \approx 1/\text{cm}^3$ is the average galactic number density of protons. Hence the life-time of “volume-annihilated” objects would be

$$\tau_{vol} = (\sigma_{ann} v n_p)^{-1} \approx 10^{15} \text{ s} \left(\frac{\text{cm}^{-3}}{n_p} \right) \left(\frac{10^{-15} \text{ cm}^3/\text{s}}{\sigma_{ann} v} \right). \quad (23)$$

The result does not depend upon the size and density of the anti-object.

Thus the low density anti-objects which are annihilated by surrounding protons inside their whole volume could not survive in galaxies to the present time, $t_U \approx 3 \cdot 10^{17} \text{ s}$. However, as we argued in Sec. 3, B-balls could naturally populate galactic halos, where the proton number density is lower, about $10^{-4}/\text{cm}^3$, and the life-time of the anti-object would be at the level of $\tau_{vol} \approx 10^{19} \text{ s}$.

The luminosity in gamma-rays of the volume-annihilated objects can be estimated as follows. The total energy release per unit of time is given by $L_{tot}^{(vol)} = 2m_p \dot{N}_{ann}$, where m_p is the proton mass. The luminosity in gamma rays, produced by the annihilation, would be

$$L_{\gamma}^{(vol)} \approx 0.3 L_{tot}^{(vol)} \approx 10^{35} \text{ erg/s} \left(\frac{\sigma_{ann} v}{10^{-15} \text{ cm}^3/\text{s}} \right) \left(\frac{R_B}{0.1 \text{ pc}} \right)^3 \left(\frac{n_p}{10^{-4} \text{ cm}^{-3}} \right) \left(\frac{n_{\bar{p}}}{10^4 \text{ cm}^{-3}} \right), \quad (24)$$

if the mean free path of photons inside the anti-object is larger than its size, R_B . If such an object is at the distance of 10 kpc from the Earth, the expected flux would be about 10^{-11} erg/s/cm², which corresponds to 10^{-7} photons/s/cm². It is below the gamma ray background with energy ~ 100 MeV and in the sensitivity range of the present and future experiments.

We would expect approximately of the same photon flux from e^+e^- -annihilation because σv for e^+e^- is twice bigger than that for $\bar{p}p$, but number of photons per annihilation is twice smaller. Thus the flux of 0.511 MeV line from an anticloud at 10 kpc would be about 10^{-7} photons/cm²/s and it should be compared with the observed flux of the same line equal to $\sim 10^{-4}$ photons/cm²/s [5].

Such anticlouds could also be observed in visible light, created by the bremsstrahlung of the energetic electrons originated from $\bar{p}p$ -annihilation. The probability of such processes is two orders of magnitude smaller than of the main one. The energy release is roughly 8 order of magnitude smaller. So the visible luminosity would be 8 orders of magnitude weaker than the solar one.

The influx of protons could be diminished by the radiation pressure, allowing the anti-objects to survive up to the present time in more dense regions as well. This effect is considered in Subsection 4.3, where we see, however, that the radiation pressure is usually not essential.

4.2 Surface annihilation

If the proton mean free path is much smaller than the size of the anti-object, $\lambda_{free} \ll R$, the annihilation takes place on the surface. This is typical situation for stellar types anti-objects and, even more, for compact anti-stars, as e.g. white dwarfs, (anti)-neutron stars, etc. In this case all the protons that hit the surface of the anti-object annihilate. The annihilation cross section is given by the geometrical area of the anti-object, $\sigma = 4\pi R^2$, and the gamma-ray luminosity of such a compact anti-object is equal to:

$$L_{\gamma}^{(sur)} \approx 4\pi R_B^2 0.6m_p F_p \approx 10^{27} \text{ erg/s} \left(\frac{R_B}{R_{\odot}} \right)^2 \left(\frac{n_p}{\text{cm}^{-3}} \right) \left(\frac{v}{10^{-3}} \right), \quad (25)$$

where $F \sim n_p v$ is the proton flux and $R_{\odot} \sim 7 \cdot 10^{10}$ cm is the Solar radius. With this luminosity a solar mass anti-star would have the life-time of the order of 10^{27} s, if all the factors in Eq. (25) are of order unity.

If such an anti-star lives in the galactic center, where $n_p \gg 1/\text{cm}^3$, its luminosity may be quite large. However, the pressure of emitted gamma radiation could reduce the proton flux and diminish $L_{\gamma}^{(sur)}$. This effect is considered in the following subsection and we see that quite high luminosities are permitted.

4.3 Eddington limit

If the luminosity of an object is created by the influx of particles, there exists an upper bound on its luminosity, which follows from the fact that the pressure of the emitted radiation diminishes the incoming flux.

The force acting on proton, or time-derivative of the proton momentum, $\dot{P}$, created by the gamma ray pressure at distance r from the radiating objects is equal to:

$$\dot{P} \sim \sigma_{p\gamma} n_{\gamma}(r) \omega, \quad (26)$$

where $\sigma_{p\gamma} \sim 10^{-31} \text{ cm}^2$ is the cross-section of the Compton scattering on proton, ω is the gamma ray energy and n_γ is the number density of emitted photons. The total luminosity of the object is $L = 4\pi R^2 n_\gamma(R) \omega$. Here n_γ is taken at the surface of the object, but it drops as $(R/r)^2$ with the increasing distance r .

The force of the gravitational attraction acting on the protons at distance r is equal to

$$F_{grav} = \frac{G_N m_p M}{r^2}. \quad (27)$$

Demanding that the gravitational attraction should be stronger than the radiation pressure, we obtain:

$$L < 10^{45} \text{ erg/s} \left(\frac{M_B}{M_\odot} \right) \left(\frac{10^{-31} \text{ cm}^2}{\sigma_{p\gamma}} \right). \quad (28)$$

One may argue that the total flux of the particles to the anti-object must be electrically neutral and hence the much larger force exerted by photons on electrons should be substituted into Eq. (26). For the low energy photons, $\omega < m_e$, the Thomson cross-section, $\sigma_{Th} = 0.66 \cdot 10^{-24} \text{ cm}^2$, should be used. In the case of photons originating from the $\bar{p}p$ -annihilation, their energies are much larger and the cross-section is about $\sigma_{e\gamma} \sim \pi \alpha^2 / (m_e \omega)$. This would diminish bound (28) by roughly 4 orders of magnitude. In any case, the Edington limit is well above prediction (25).

On the other hand, the excessive charge created by a larger influx of protons with respect to electrons could be compensated by the out-flux of positrons from the antimatter object. If this is the case, the limit on the gamma-ray luminosity would be given by Eq. (28). Energetic gamma rays from $\bar{p}p$ -annihilation would be accompanied by the emission of low energy positrons, which would be a source of 0.511 MeV line.

5 Point-like sources of gamma radiation

First, we consider the possibility of presence of anti-clouds in our galaxy. If condition (21) is satisfied, the proton mean free path inside the anti-cloud is larger than the anti-cloud size and matter-antimatter annihilation proceeds in the whole volume. The life-time of such clouds is given by Eq. (23), if the proton flux is sufficiently large. Such clouds would not survive to the present time. However, if the proton flux is not big enough, the life-time of the cloud may exceed the age of the Galaxy.

More favorable condition for survival of anti-clouds are in the galactic halo, where the proton density is $n_p \sim 10^{-4} \text{ cm}^{-3}$. The gamma-ray luminosity of such a cloud is given by Eq. (24). Such a source might be observed on the Earth as a γ ray source with the flux:

$$\phi_{Earth} \sim 10^{-7} \left(\frac{n_p}{10^{-4} \text{ cm}^{-3}} \right) \left(\frac{n_{\bar{p}}}{10^4 \text{ cm}^{-3}} \right) \left(\frac{R_B}{0.1 \text{ pc}} \right)^3 \left(\frac{10 \text{ kpc}}{d} \right)^2 \text{ cm}^{-2} \text{ s}^{-1}, \quad (29)$$

where d is the distance of the anti-cloud from the Earth. Eq. (29) should be compared with the point source sensitivity of EGRET [24], at the level of $10^{-7} \text{ photons cm}^{-2} \text{ s}^{-1}$ for $E_\gamma > 100 \text{ MeV}$ and a full two weeks exposure, and of the near-future GLAST [25], which is about two order of magnitude better, $\sim 10^{-9} \text{ photons cm}^{-2} \text{ s}^{-1}$.

As for possible anti-stars, emitting gamma rays from $p\bar{p}$ -annihilation on their surface, they should be quite close to us in order to be detectable point-like sources. For an anti-star

in the galactic disc, its γ flux would be

$$\phi_{Earth} \sim 10^{-7} \left(\frac{R_B}{R_\odot} \right)^2 \left(\frac{1 \text{ pc}}{d} \right)^2 \text{ cm}^{-2} \text{ s}^{-1}. \quad (30)$$

To be observable in a near future such an anti-star should be really in solar neighborhood and if it is a normal star powered by thermonuclear energy it could even be seen with a naked eye. On the other hand, if anti-stars are similar to the Sun, the consideration of a possible anti-stellar wind requires that their number in the Galaxy is smaller than 10^5 (see below Eq. (38)) and, assuming they are uniformly distributed in the galactic disk, we would expect a mean anti-star number density of $1/(100 \text{ pc})^3$. In this case $d \sim 100 \text{ pc}$ implies $\phi_{Earth} \sim 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$, a photon flux too weak to be detectable from a point-like source.

The γ -flux from an anti-star may be strongly enhanced if it happens to be in a high density hydrogen cloud. This possibility is quite realistic: even if the mean proton density is $n_p \sim 1 \text{ cm}^{-3}$, about one half of the interstellar medium is tied up in gas clouds with average proton density $n_p \sim 10^3 \text{ cm}^{-3}$. In other words, gas clouds occupy a fraction of about 10^{-3} of the Galaxy volume, so that, assuming 10^5 anti-stars in the Galaxy, about 100 of them are expected to live in some gas cloud. Their gamma flux would increase by three orders of magnitude, or even more if we consider that some clouds are much denser, up to $n_p \sim 10^6 - 10^9 \text{ cm}^{-3}$.

Last, it could be interesting to consider observational signatures of anti-black holes, i.e. black holes generated by the gravitational collapse of antimatter. They may be distinguished from the ordinary black holes if they were surrounded by an atmosphere of antimatter (it is not difficult to imagine that it is possible, even if detailed calculations would be necessary for a precise assertion). Their anti-atmosphere could be considered as an anti-cloud around a very compact anti-object and, following the considerations of Subsection 4.1, such an anti-atmosphere could survive up to the present day only if such an anti-black hole was in the galactic halo. In this case this strange anti-object could be detectable looking for high energy γ radiation (see Eq. (29)) from a stellar mass object creating gravitational micro-lensing (MACHO) [26]. The same would be true also for compact anti-stars in galactic halo.

The spectrum of photons from matter-antimatter annihilations is well known and consists of three different parts. First, most energetic are photons from $\bar{p}p \rightarrow \pi^0 \rightarrow 2\gamma$. If π^0 were at rest we would observe the single 67.5-MeV line. However, the life-time of π^0 is very short and they decay being relativistic. Thus the spectrum spreads both ways up and down and shifts from zero to higher energies above 200 MeV. The second less energetic and also continuous part comes from the chain of reactions $\bar{p}p \rightarrow \pi^+ \rightarrow \mu^+ \rightarrow e^+$ and subsequent e^+e^- -annihilation. The probability of such double annihilation and the shape of the spectrum depends upon the object where these processes take place. This process seems to be least efficient. The annihilation of the slow positrons inside antimatter objects with the background electrons leads to the famous 0.511 MeV line, which is easy to identify. This anomalously bright line is observed recently in the Galactic center [27], Galactic bulge [4] and possibly even in the halo [5]. Though an excess of slow positrons is explained in a conventional way as a result of their creation by light dark matter particles, such a suggestion is rather unnatural because it requires a fine-tuning of the mass of the dark matter particle and the electron mass. More natural explanation is the origin of these positrons from primordial antimatter objects.

6 Diffuse galactic gamma ray background

In the standard theory, the galactic production of γ rays is due to inelastic collisions of high energy cosmic rays with the interstellar medium (the dominant processes are $p + p$, $p + \alpha$ and $\alpha + \alpha$), to bremsstrahlung radiation from cosmic ray electrons and to inverse Compton scattering of electrons with low energy photons. Many authors have calculated the galactic production rate of γ per hydrogen atom (see e.g. Ref. [28]) and the estimated rate in the energy range $E_\gamma > 100$ MeV is

$$\Gamma_\gamma \sim 2 \cdot 10^{-25} \text{ s}^{-1} \text{ atom}^{-1}, \quad (31)$$

in good agreement with observational data [29]. From Eq. (31) we can deduce the total production rate of high energy γ rays in our galaxy

$$\Gamma_\gamma^{tot} \sim 10^{43} \text{ s}^{-1}. \quad (32)$$

The presence of $N_{\bar{S}}$ anti-stars in the galactic disc, where the average proton number density is $n_p \sim 1 \text{ cm}^{-3}$, would create a new source of high energy γ rays, with the contribution

$$\dot{N}_\gamma \sim \dot{N}_\gamma^{(sur)} N_{\bar{S}}, \quad (33)$$

where $\dot{N}_\gamma^{(sur)} \approx L_\gamma^{(sur)}/(100 \text{ MeV})$ is the photon flux coming from each anti-star and $L_\gamma^{(sur)}$ is given by Eq. (25). If we assume that $\dot{N}_\gamma$ (33) cannot exceed 10% of the standard galactic production rate of high energy γ , given by Eq. (32), we obtain a bound on the present number of anti-stars (for simplicity, we assume that all the anti-stars have the same radius R_B)

$$N_{\bar{S}} \lesssim 10^{12} \left(\frac{R_\odot}{R_B} \right)^2. \quad (34)$$

A stronger constraint can be obtained from the consideration of the annihilation of antimatter from the anti-stellar wind with protons in the interstellar medium. The rate of $\bar{p}$ emission by the anti-stellar wind per anti-star is

$$\dot{N}_{\bar{p}}^{wind} \sim 10^{36} W \text{ s}^{-1}, \quad (35)$$

where $W = \dot{M}/\dot{M}_\odot$ is the ratio of an anti-star mass loss rate to the Solar one. The total number of antiprotons in the Galaxy can be determined by an equation:

$$\dot{N}_{\bar{p}}^{tot} = S - T, \quad (36)$$

where $S = \dot{N}_{\bar{p}}^{wind} N_{\bar{S}}$ is the source term and T the sink term due to annihilation of $\bar{p}$. The life-time of antiprotons in the Galaxy with respect to annihilation is given by Eq. (23), i.e. it is much smaller than the age of the Galaxy. Thus the number of antiprotons reached a stationary value, i.e. $\dot{N}_{\bar{p}}^{tot} = 0$ and the production rate of 100 MeV γ in the Galaxy is $\dot{N}_\gamma \approx 4S$, because in each act of $\bar{p}p$ -annihilation 4 photons are produced on the average. The flux of 100 MeV photons on the Earth would be

$$\phi_{\bar{p}p}^{(gal)} \approx \frac{4\dot{N}_{\bar{p}}^{wind} N_{\bar{S}}}{4\pi R_{gal}^2} \approx 100W \left(\frac{N_{\bar{S}}}{N_S} \right) \text{ cm}^{-2}\text{s}^{-1}, \quad (37)$$

where we took $N_S = 3 \cdot 10^{11}$ and $R_{gal} = 10$ kpc. The luminosity of the Galaxy in 100 MeV gamma rays from anti-stellar wind would be $L_{\bar{S}} \sim 10^{44} W N_{\bar{S}}/N_S$ erg/s. Since from Eq. (32) we find that the total Galaxy luminosity in 100 MeV γ is $L_{\gamma}^{tot} \sim 10^{39}$ erg/s, the related bound on the number of anti-stars is:

$$N_{\bar{S}}/N_S \lesssim 10^{-6} W^{-1}. \quad (38)$$

Here, as always we assume that the contribution from new physics cannot exceed 10% of L_{γ}^{tot} .

A similar restriction can also be obtained from the 0.511 MeV line created by e^+e^- annihilation with positrons from the anti-stellar wind. Since the number of antiprotons in the stellar wind is approximately the same as the number of positrons, the flux of 0.511 MeV photons would be close to that given by Eq. (37). Taking that the latter is smaller than 10^{-4} /cm²/s we find

$$N_{\bar{S}}/N_S \lesssim 10^{-6} W^{-1}. \quad (39)$$

If anti-stars have been formed in the very early universe in the regions with a high antibaryonic density [17], such primordial stars would most probably be compact ones, white or brown dwarfs, neutron stars, etc. The stellar wind in this case would be much smaller than the solar one, $W \ll 1$. Their luminosity from the annihilation on the surface should be very low, because of their small radius R , and their number in the Galaxy may be as large as the number of the usual stars. This possibility is not excluded by the bounds (34) and (38). Such compact dark stars could make a noticeable part of the cosmological dark matter. As we have argued in Sec. 3, the early created compact stellar like objects behave as the usual collisionless cold dark matter (CDM). In this case it is natural to expect that they would be distributed in and around galaxies as the standard CDM, having a large number density in the galactic center and decreasing as $1/r^2$ in the halo.

These compact objects would generate the diffuse gamma-ray background not only now but during all cosmological history. In particular, we expect a strong constraint on the number of anti-stars from their emitted radiation during the so called “dark age”, after recombination but before the advent of the early standard stars. We thank Francesco Villante for having pointed out this possibility. We plan to evaluate the intensity of such cosmological γ -background in another work.

7 Anti-nuclei in cosmic rays

Stable charged particles in cosmic rays consist of 86% of protons, 11% helium nuclei, 1% heavier nuclei, and 2% electrons. It is common belief that the abundances of the elements in the cosmic rays reflect relative abundances in the Galaxy (even if the low energy cosmic rays should be a mirror of the relative abundances in the Solar System). Hence, we can reasonably expect that the antimatter-matter ratio in cosmic rays is more or less equal to the (anti-star)-star ratio $N_{\bar{S}}/N_S$, if the antistars are of the same kind as the stars in the Galaxy. As for antiprotons and positrons, they are not direct indicators for the existence of primordial antimatter, because they can be produced in many astrophysical processes. For example, the observed $\bar{p}/p$ ratio is at the level of 10^{-4} and is compatible with theoretical predictions for $\bar{p}$ production by the high energy cosmic ray collisions with the interstellar medium. A possible contribution of $\bar{p}$ from exotic sources (ES) is not more than about 10%

of the total observed $\bar{p}$ flux, so the number of anti-stars $N_{\bar{S}}$ have to be

$$\frac{N_{\bar{S}}}{N_S} \lesssim \left(\frac{\bar{p}}{p} \right)_{\text{ES}} \lesssim 10^{-5} \quad \Rightarrow \quad N_{\bar{S}} \lesssim 10^6, \quad (40)$$

since the number of ordinary stars in the Galaxy is $N_S \sim 10^{11}$.

On the other hand, the possibility of producing heavier anti-nuclei (such as anti-helium) in cosmic ray collisions is completely negligible and a possible future detection of the latter would be a clear signature of antimatter objects. At present there exists an upper limit on the anti-helium to helium ratio in cosmic rays, at the level of 10^{-6} [30], leading to the constraint

$$\frac{N_{\bar{S}}}{N_S} \lesssim \left(\frac{\bar{He}}{He} \right)_{\text{ES}} \lesssim 10^{-6} \quad \Rightarrow \quad N_{\bar{S}} \lesssim 10^5, \quad (41)$$

which is essentially equal to that from anti-stellar wind in Eq. (38). However, the sensitivity of AMS [31] and PAMELA [32] space missions is expected to be two orders of magnitude better, at about 10^{-8} . In this case the non-observation of anti-helium nuclei would lead to the much stronger constraint

$$N_{\bar{S}} \lesssim 10^3. \quad (42)$$

Of course, the bound (42) cannot be applied if anti-stars are compact ones from the very beginning (i.e. from the moment of their formation in the early universe). In this case the stellar wind from them and the shortage of anti-supernova events would spread much less (anti)helium than the normal stars, but heavier anti-elements might be not so strongly suppressed. According to the scenario described in Sec. 3, the compact (anti-)stars might form in (anti)baryon-rich regions, where the primordial nuclear abundances would be quite different from the standard ones with an enhanced amount of heavier (anti-)nuclei, as e.g. oxygen, nitrogen, carbon, and maybe even heavier one up to calcium or iron [21]. Spectroscopy observations may detect stellar atmospheres anomalously rich in heavy elements. Such chemical anomaly is a good signature that such object are made of antimatter and a search for gamma rays from them looks promising. On the other hand, the mechanisms capable of ejecting antimatter from such stars and of spreading it out in the Galaxy are favorable for enrichment of galactic cosmic rays with heavier anti-nuclei. Unfortunately the amount of anti-nuclei depends upon many unknowns and it is impossible to make a reliable estimate of their flux.

8 More exotic events

The presence of anti-stars in the Galaxy could lead to extraordinary events of star-antistar annihilation. As a matter of fact, the radiation pressure produced in the collision prevents their total destruction. Still the released energy can be huge.

The most spectacular phenomenon is a collision between a star and an anti-star with similar masses M . The energy released in such a collision can be estimated as follows. The relative momentum of the colliding stars is approximately $P \sim Mv$, where the typical value of the relative velocity is about $v \approx 10^{-3}$. This momentum would be “canceled” by the pressure of radiation created by baryon-antibaryon annihilation, p . The total pressure force is $F \sim \pi r^2 p$, where r is the radius of the circle where the colliding stars penetrated into each

other. The bounce would take place when $Ft_{coll} \sim Mv$, where $t_{coll} \sim d/v$ is the collision time and d is the penetration depth. Since the annihilation products are relativistic, their pressure density is of the same order of magnitude as the energy density and hence the amount of the annihilated matter during the star-antistar collision would be

$$m_{ann} \sim Mv^2, \quad (43)$$

i.e. the total energy release would be

$$\Delta E_1 \sim 10^{48} \text{ erg} (M/M_\odot)(v/10^{-3})^2. \quad (44)$$

Most probably the radiation would be emitted in a narrow disk along the boundary of the colliding stars.

The collision time can be estimated as follows. Let us introduce the angle θ at which the radius, r , of the collision disk is seen from the star center, $r = R\theta$, assuming that $\theta < 1$. Here R is the star radius. The penetration depth is $d = R\theta^2$. The volume where matter and antimatter mix and annihilate is $R^3\theta^4$. This volume should be about v^2 of the total stellar volume. Thus the penetration depth is $d \sim vR$ and the collision time is $t_{coll} \sim R$. For the solar type star this time is about 3 s. For colliding compact star-antistar the collision time would be much smaller but the energy release could be much larger, because the velocity might approach relativistic values.

We expect that in the process of a star collision with the similar anti-star the energy would be emitted inside a narrow disk with the opening angle $\theta \sim \sqrt{v}$. The characteristic time of the emission is of the order of a second. The energy of the radiation should be noticeably smaller than 100 MeV because the radiation should degrade in the process of forcing the star bounce. This makes this collision similar to gamma bursts but unfortunately some other features do not fit so well: the released energy should be much larger, about $10^{53} \sqrt{v}$ erg and it is difficult to explain the features of the afterglow. A reasonable amount of the energy could be released in the case of compact star annihilation. It may be possible to explain two or more bursts by the oscillating motion with interchangingly dominated attraction by gravity and repulsion by the pressure of the products of annihilation. The process is surely more complicated than our naive picture and we cannot exclude that gamma bursts are the results of star-antistar annihilation.

The collision of a compact star, e.g. an anti-neutron star, with a usual one or with a red giant would look completely different. In this case, the mass densities of the two objects are so much diverse that the anti-neutron star would go through the red giant without stopping and would annihilate all what it meets on the way. The released energy is about

$$\Delta E_2 \sim \pi R_{ns}^2 D \rho_{rg} = M_{rg} \left(\frac{R_{ns}}{R_{rg}} \right)^2 \left(\frac{D}{R_{rg}} \right), \quad (45)$$

where R_{ns} is the radius of the anti-neutron star, D is the crossed distance inside the red giant, R_{rg} is the radius of the red giant, ρ_{rg} is the red giant mass density, and M_{rg} is its mass. For $M_{rg} \sim M_\odot$ and $R_{rg} \sim 10^{14}$ cm a reasonable estimate of this energy release is 10^{38} erg. The crossing time of the red giant is about $R_{rg}/v \sim 3 \cdot 10^6$ seconds. Hence the additional luminosity during anti-neutron star propagation inside the red giant would be an order of magnitude smaller than the solar luminosity and, probably, most of the energy would reheat the interior of the star and could not reach the free space. These estimates are valid for collision of a compact antistar with the envelope of red giant. A collision of

antistar with the core of red giant would look similar to the compact star-antistar collision considered above. Most spectacular would be a collision of red-giant with anti-red-giant, where a fantastic amount of energy would be released.

The probability of the collision of two stars can be estimated as follows. The collision rate is $\Gamma = \sigma n_S v$, where $\sigma \approx \pi R^2$ is the geometrical area of the larger star and $n_S \sim 1/\text{pc}^3$ the mean star number density in the Galaxy. The total number of collisions per unit time would be:

$$\dot{N}_{\bar{S}} = \Gamma N_{\bar{S}} = \pi v R^2 N_{\bar{S}} n_S \approx 10^{13} \text{ year}^{-1} \left(\frac{N_{\bar{S}}}{10^5} \right) \left(\frac{R}{10^{11} \text{ cm}} \right)^2. \quad (46)$$

If anti-stars are similar to the Sun, from the bound (38) we find $N_{\bar{S}} \lesssim 10^5$, which implies essentially no collisions during the whole history of the universe. On the other hand, if the anti-stars are white dwarfs or anti-neutron stars, the bounds (34) and (38) are very weak or inapplicable and the anti-star number can be as large as that of the ordinary stars, i.e. $N_{\bar{S}} \sim 10^{11}$. In this case we find one collision per 10^7 years.

The collision with a red giant would have a larger cross-section because of much larger radius of the red giant, $R_{rg} > 10^{13}$ cm. Since red giants are about 1% of the stars in the Galaxy, the number of collisions could increase by 2 orders of magnitude. Another factor which might also enhance the probability is a larger gravitational attraction of heavy stars. We know, the majority of the stars in the Galaxy are in multiple stellar systems. However, it is surely true for the normal stars which were formed from the primeval hydrogen cloud in galactic disk, while B-balls were created in the early universe by a completely different mechanism. However, the gravitational capture of a normal star and B-ball has the same probability as the gravitational capture of the normal stars, see discussion in the next paragraph.

Another interesting possibility is the transfer of material in a binary system. In the case of the ordinary stars made of matter, it can happen that a binary system is formed by a red giant with a close compact star companion such as a white dwarf. If the former overflows the Roche lobe of the latter, the compact star captures gases from the red giant outer atmosphere. Then, on the white dwarf surface, a large amount of hydrogen is rapidly converted into heavier elements via the CNO cycle, producing an extremely bright outburst of light. The event taking place can be a nova, where the star luminosity at the brightness peak is about 10^{38} erg/s, or the much more spectacular phenomenon of supernova Ia, with a maximum luminosity of about 10^{52} erg/s (however, the latter case is not of interest for us, because the emitted energy comes from star gravitational collapse and does not from the CNO cycle). We can reasonably expect that something similar can happen in a binary system where one of the stars is made of matter and the other one of antimatter. In this case hydrogen is not burnt via the CNO cycle, but the much more violent process of matter-antimatter annihilation takes place and the white dwarf should be brighter than an ordinary nova. Of course, the two stars cannot be born at the same time from the same cloud, but they must have a different origin. For example, one star can be gravitationally captured by another which is already in a multiple star system (energy conservation forbids the formation of bound systems from two stars) and, even if today it is ruled out as the dominant formation mechanism, it is expected to be not rare (see e.g. Ref. [33]). The event could be observed as a long outburst of 100 MeV γ radiation and be easily distinguished from the standard novae: the star luminosity could not exceed the Eddington limit (28), but would be probably close to it. Moreover, even if the rate of the ordinary novae in the Galaxy is estimated to be only about 50 events per year, thanks to the large amount of

energy release, we can monitor many galaxies at the same time, even quite distant from us, and increase our possibility of observing this kind of phenomena. Even in this case, clear predictions are impossible, because observational probabilities depend on many unknowns.

9 Conclusion

The conclusion to this paper, unfortunately, is not conclusive – practically anything is allowed. Gamma rays from $\bar{p}p$ -annihilation may be observable with future or even with existing γ -telescopes. Quite promising for discovery of cosmic antimatter are point-like sources of gamma radiation. The problem is to identify a source which is suspicious to consist of antimatter. A possible manifestation of such a source is an anomalous abundance of chemical (anti-)elements around it, which can be measured by spectroscopy.

The 100-MeV gamma ray background does not have pronounced features which would unambiguously tell that the photons came from the annihilation of antimatter. The photons produced as a result of $\bar{p}p$ annihilation would have a well known spectrum but it may be difficult to establish a small variation of the conventional spectrum due to such photons.

In contrast, the 0.511 MeV line must originate from e^+e^- annihilation and it is tempting to conclude that the observed excessive signal from the Galaxy and, especially, from the galactic bulge come from astronomical antimatter objects.

An important feature of the scenario of early formed compact antimatter objects [17] is that such stellar type ones would behave as normal cold dark matter and they would concentrate in or around normal galaxies.

Interesting candidates for being anti-matter stars are the observed MACHO events [26]. According to the scenario of Ref. [17], these stellar mass gravitational lenses should consist of practically equal number of matter and antimatter objects. The latter should emit 100 MeV and 0.511 MeV gamma rays and, though the luminosity might be rather weak, see Eq. (25), they still may be observed with high angular resolution telescopes. The gamma ray luminosity would be much stronger for MACHOs in the galactic disk, because the number density of protons there is at least 4 orders of magnitude higher than in the galactic halo.

If an antistar happens to be in the Galactic Center, its luminosity from the surface annihilation of the background matter should be strongly enhanced due to the much larger density of the interstellar matter there. So the search of the antimatter signatures in the direction of the Center is quite promising.

There is a non-negligible chance to detect cosmic anti-nuclei and not only light anti-helium but also much heavier ones, especially if anti-stars became early supernovae.

A possible discovery of cosmic antimatter would shed light on the mechanism of baryogenesis and CP-violation in cosmology. In the standard scenarios the baryon asymmetry β is constant, just one number, and it is impossible to distinguish between different mechanisms of baryogenesis measuring this single number. More exotic models predicting a noticeable amount of antimatter in our neighborhood are much more interesting because $\beta = \beta(x)$ is a function of space points and contains much more information about physics in the early universe.

Macroscopically large pieces of antimatter not far from the Earth may be interesting energy sources, but this is not in foreseeable future, maybe only in science fiction.

Acknowledgments

We thank Filippo Frontera and Francesco Villante for useful comments and suggestions.

References

- [1] G. Steigman, *Ann. Rev. Astron. Astrophys.* **14**, 339 (1976).
- [2] F.W. Stecker, *Nucl. Phys. B* **252**, 25 (1985).
- [3] M. Boezio *et al.* (WIZARD Collaboration), *Astrophys. J.* **487**, 415 (1997);
S. Orito *et al.* (BESS Collaboration), *Phys. Rev. Lett.* **84**, 1078 (2000),
astro-ph/9906426.
- [4] G. Weidenspointner *et al.*, *Astron. Astrophys.* **450**, 1013 (2006), astro-ph/0601673.
- [5] G. Weidenspointner *et al.*, astro-ph/0702621.
- [6] W.-M. Yao *et al.*, *Journal of Physics G* **33** (2006) 1. <http://pdg.lbl.gov/>.
- [7] D.N. Spergel *et al.*, *Astrophys. J. Suppl.* **148**, 175 (2003), astro-ph/0302209.
- [8] Ya.B. Zeldovich, *ZhETF*, **48**, 986 (1965);
A.D. Dolgov and Ya.B. Zeldovich, *Rev. Mod. Phys.* **53**, 1 (1981);
E.W. Kolb and M.S. Turner, *The Early Universe* (Addison-Wesley, Redwood City, USA, 1990).
- [9] A.D. Dolgov, *Phys. Repts.* **222**, 309 (1992).
- [10] A.D. Dolgov, hep-ph/9707419, lectures at 25th ITEP Winter School of Physics, Moscow, Russia, 18 – 27 February, 1997.
- [11] A. Riotto and M. Trodden, *Ann. Rev. Nucl. Part. Sci.* **49**, 35 (1999), hep-ph/9901362.
- [12] V.A. Rubakov and M.E. Shaposhnikov, *Phys. Usp.* **39**, 461 (1996); *Usp. Fiz. Nauk* **166**, 493 (1996), hep-ph/9603208.
- [13] A.G. Cohen, A. De Rújula and S.L. Glashow, *Astrophys. J.* **495**, 539 (1998),
astro-ph/9707087;
A.G. Cohen and A. De Rújula, astro-ph/9709132.
- [14] K.M. Belotsky, Yu.A. Golubkov, M.Yu. Khlopov, R.V. Konoplich and A.S. Sakharov, *Phys. Atom. Nucl.* **63**, 233 (2000), *Yad. Fiz.* **63**, 290 (2000), astro-ph/9807027;
Yu.A. Golubkov and M.Yu. Khlopov, *Phys. Atom. Nucl.* **64**, 1821 (2001), *Yad. Fiz.* **64**, 1904 (2001), astro-ph/0005419;
D. Fargion and M. Khlopov, *Astropart. Phys.* **19**, 441 (2003), hep-ph/0109133.
- [15] A.D. Dolgov, hep-ph/0511213, lectures presented at Varenna School *CP Violation: From Quarks to Leptons*, Varenna, Italy, July, 2005.
- [16] A.D. Dolgov, Antihydrogen Workshop, Munich, Germany, July, 1992, hep-ph/9211241;
A.D. Dolgov, *Nucl. Phys. Proc. Suppl.* **113**, 40 (2002);
A.D. Dolgov, in *Proceedings of 14th Rencontres de Blois: Matter - Anti-matter Asymmetry*, Chateau de Blois, France, 17-22 June, 2002, hep-ph/0211260.

- [17] A. Dolgov and J. Silk, Phys. Rev. D **47**, 4244 (1993).
- [18] I. Affleck and M. Dine, Nucl. Phys. B **249**, 361 (1985).
- [19] A. Vilenkin and L. Ford, Phys. Rev. D **26**, 1231 (1982);
A.D. Linde, Phys. Lett. B **116**, 335 (1982);
A.A. Starobinsky, Phys. Lett. B **117**, 175 (1982);
A.D. Dolgov and D.N. Pelliccia, Nucl. Phys. B **734**, 208 (2006).
- [20] R.B. Metcalf and J. Silk, astro-ph/0612253.
- [21] S. Matsuura, A.D. Dolgov, S. Nagataki and K. Sato, Prog. Theor. Phys. **112**, 971 (2004);
S. Matsuura, S.I. Fujimoto, S. Nishimura, M.A. Hashimoto and K. Sato, Phys. Rev. D **72**, 123505 (2005), astro-ph/0507439.
- [22] D.H. Oaknin and A.R. Zhitnitsky, Phys. Rev. D **71**, 023519 (2005), hep-ph/0309086;
A.R. Zhitnitsky, JCAP 10 (2003) 010, hep-ph/0202161;
- [23] D.H. Oaknin and A.R. Zhitnitsky, Phys. Rev. Lett. **94**, 101301 (2005), hep-ph/0406146;
K. Lawson and A.R. Zhitnitsky, arXiv:0704.3064.
- [24] <http://heasarc.gsfc.nasa.gov/docs/cgro/egret/>
- [25] <http://glast.stanford.edu/>
- [26] C. Alcock *et al.*, Astrophys. J. **486**, 697 (1997), astro-ph/9606165.
- [27] P. Jean *et al.*, Astron. Astrophys. **407**, L55 (2003), astro-ph/0309484;
J. Knödseder *et al.*, Astron. Astrophys. **411**, L457 (2003), astro-ph/0309442;
J. Knödseder *et al.*, Astron. Astrophys. **441**, 513 (2005), astro-ph/0506026.
- [28] M. Mori, Astrophys. J. **478**, 225 (1997), astro-ph/9611233.
- [29] S.W. Digel, S.D. Hunter and R. Mukherjee, Astrophys. J. **441**, 270 (1995).
- [30] M Sasaki *et al.*, Nucl. Phys. B Proc. Suppl. **113**, 202 (2002).
- [31] <http://ams.cern.ch/>
- [32] <http://wizard.roma2.infn.it/pamela/>
- [33] C. Clarke, in *The Formation of Binary Stars*, IAU Symposium **200** (2001), eds. H. Zinnecker and R.D. Mathieu.